

Towards a Nuclear Charge Radius Determination of the One-Neutron Halo Nucleus ^{11}Be

M. Žáková^a, D. Tiedemann^a, C. Geppert^{a,b,c}, R. Sánchez^b, Z. Andjelkovic^a,
K. Blaum^{b,d}, M.L. Bissell^e, G.W.F. Drake^f, J. Eble^g, A. Herlert^h,
M. Kowalska^h, J. Krämer^a, A. Krieger^a, R. Neugart^d, F. Schmidt-Kaler^g,
Z.-C. Yanⁱ, D. Yordanov^d, C. Zimmermann^c, and W. Nörtershäuser^{a,b}

^a*Institut für Kernchemie, Universität Mainz, 55099 Mainz, Germany*

^b*Gesellschaft für Schwerionenforschung, 64291 Darmstadt, Germany*

^c*Institut für Physik, Universität Tübingen, 72076 Tübingen, Germany*

^d*Institut für Physik, Universität Mainz, 55099 Mainz, Germany*

^e*Instituut voor Kern- en Stralingsfysica, KU Leuven, 3001 Leuven, Belgium*

^f*Department of Physics, University of Windsor, Windsor, ON, Canada, N9B 3P4*

^g*Institut für Quanteninformationsverarbeitung, Universität Ulm, 89081 Ulm, Germany*

^h*CERN, CH-1211 Geneva 23, Switzerland*

ⁱ*Department of Physics, University of New Brunswick, Fredericton, NB, Canada E3B 5A3*

Using very accurate isotope shift measurements combined with mass shift calculations at the highest level of accuracy that is nowadays available for two- and three-electron systems, nuclear charge radii of the lightest elements can be extracted in a nuclear-model independent way [1]. Such measurements are challenging because in light nuclei, the nuclear volume effect of the isotope shift is a tiny fraction (10^6 Hz) of the total isotope shift (3×10^{10} Hz) in an optical transition (10^{15} Hz). Particular interesting is the application of this method for exotic short-lived isotopes of helium and lithium, namely the two-neutron halo nuclei ^6He [2], ^{11}Li [3], and the four-neutron halo nucleus ^8He [4]. In extension of these studies on light nuclides close to the neutron-dripline, a first calculation of the mass shift in Be^+ has been completed [5], and will be used to determine the charge radius of the one-neutron halo nucleus ^{11}Be . High-accuracy isotope shift measurements of $^{7,9,10,11}\text{Be}$ are currently being prepared at the ISOLDE facility at CERN, based on high-resolution fluorescence spectroscopy involving two frequency doubled dye lasers and an optical frequency comb for stabilization and frequency determination. The setup has recently been successfully used on a test beam of the stable isotope ^9Be . The technique and the results of this first off-line experiment will be presented.

[1] G.W.F. Drake et al., Lecture Notes in Physics **745** 131 (2008).

[2] L.-B. Wang et al., Phys. Rev. Lett. **93** 142501 (2004).

[3] R. Sánchez et al., Phys. Rev. Lett. **96** 033002 (2006).

[4] P. Müller et al., Phys. Rev. Lett. **99** 252501 (2007).

[5] Z.-C. Yan et al., Phys. Rev. Lett., *in print* (2008).